

Is your child up to date on immunizations?

Vaccines can prevent or help protect against deadly and difficult-to-treat diseases. And the shots don't end after infancy. Check below for the recommended ages of each immunization and determine if your child is current with his or her vaccines. Talk with your provider about an immunization schedule for your child.

Birth

- Recombivax HB (Hepatitis B)

2 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (*Haemophilus influenzae* type B)

4 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (*Haemophilus influenzae* type B)

6 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (*Haemophilus influenzae* type B)

12 months

- M-M-R II (Measles, Mumps, Rubella)
- Varivax (Varicella)
- Prevnar 13 (Pneumococcal conjugate)
- Vqta (Hepatitis A)

15 months

- Infanrix (Diphtheria, tetanus, and acellular pertussis [DTaP])
- Hiberix (*Haemophilus influenzae* type B)*

18 months

- Vqta (Hepatitis A)

4–6 years

- ProQuad (Measles, Mumps, Rubella/Varivax [Varicella]) or separate injections
- Kinrix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus)

11–12 years

- Boostrix (Tetanus, diphtheria, and acellular pertussis [Tdap])
- Menveo (Meningococcal Group A, C, Y)
- Gardasil 9 (Human papillomavirus)—2 doses 6 months apart (over age 15 years requires 3 doses)

16 years

- Menveo (Meningococcal Group A, C, Y)
- Bexsero (Meningococcal Group B)—two doses**

Influenza

- Yearly for all children ages 6 months and up, if no contraindications

NOTE: This schedule is for preventative vaccinations. For catch up schedules or for patients with chronic conditions, see the CDC Immunization Schedule for vaccine recommendations.

*Can be administered as early as 12 months old

**For adolescents not at increased risk, age 16-23 years (preferred age 16-18 years) is based on shared clinical decision making, 2 doses 1 month apart.

