B-MEDS & PROTOCOLS

Post-Test

1. *The correct dosage of Albuterol.*

   a. *2.5mg / 3cc normal saline*
   b. *5mg / 3cc normal saline*
   c. *10mg / 3cc normal saline*
   d. *25mg / 3cc normal saline*

2. *Albuterol is areolized with:*

   a. *6 Liters O2*
   b. *2Liters O2*
   c. *15 Liters O2*
   d. *25 Liters 02*

3. *Some side effects of Albuterol are:*

   a. *Palpatations*
   b. *Dizziness*
   c. *Nervousness*
   d. *All of the above.*

4. *The correct dose of Epinephrine via auto injector is:*

   a. *0.3mg lateral thigh*
   b. *3.0mg deltoid*
   c. *1mg SL*
   d. *None of the above.*

5. *How long is the Epinephrine auto injector left in place after administration?*

   a. *30 Seconds*
   b. *60 Seconds*
   c. *2 Seconds*
   d. *10 Seconds*

6. *Oral Glucose is used <u>only</u> in an unresponsive patient.*

   a. *True*
   b. *False*

7. *The correct dose of Glucagon is:*

   a. *1mg*
   b. *.01mg*
   c. *4mg*
   d. *15 grams*

8. *Does initial Nitroglycerin require base station contact before BLS administration?*

   a. *Yes*
   b. *No*

9. *Contraindications to Aspirin include asthma, and active ulcer disease?*

   a. *True*
   b. *False*

10. *BLS may administer Nitroglycerin ___ time(s) before base station contact?*

   a. *1*
   b. *3*
   c. *None BLS must call before giving Nitroglycerin*
   d. *BLS does not give Nitroglycerin*

11. *Headache, dizziness and hypotension are side effects of Nitroglycerin?*

   a. *True*
   b. *False*

12. *Combi-Tubes require Base Station contact before insertion.*

   a. *True*
   b. *False*

13. *How much Aspirin is given to the cardiac patient?*

   a. *324mg (4 tablets of 81mg)*
   b. *1296mg (4 tablets of 324mg)*
   c. *500mg (5 tablets of 81mg)*
   d. *100mg (2 tablets of 81mg)*