B-MEDS & PROTOCOLS

Post-Test

1. *The correct dosage of Albuterol.*

    a. *2.5mg / 3cc normal saline*
    b. *5mg / 3cc normal saline*
    c. *10mg / 3cc normal saline*
    d. *25mg / 3cc normal saline*

2. *Albuterol is areolized with:*

    a. *6 Liters O2*
    b. *2Liters O2*
    c. *15 Liters O2*
    d. *25 Liters 02*

3. *Some side effects of Albuterol are:*

    a. *Palpatations*
    b. *Dizziness*
    c. *Nervousness*
    d. *All of the above.*

4. *The correct dose of Epinephrine via auto injector is:*

    a. *0.3mg lateral thigh*
    b. *3.0mg deltoid*
    c. *1mg SL*
    d. *None of the above.*

5. *How long is the Epinephrine auto injector left in place after administration?*

    a. *30 Seconds*
    b. *60 Seconds*
    c. *2 Seconds*
    d. *10 Seconds*

6. *Oral Glucose is used <u>only</u> in an unresponsive patient.*

    a. *True*
    b. *False*

7. The correct dose of Glucagon is:

a. Img
b. .0lmg
c. *4mg*
d. *15 grams*

8. Does Nitroglycerin require base station contact before BLS administration?

a. Yes
b. No

9. Contraindications to Aspirin include asthma, and active ulcer disease?

a. True
b. False

10. BLS may administer Nitroglycerin _ timers) before base station contact?

a. 1
b. 3
c. *None BLS must call before giving Nitroglycerin*
d. *BLS does not give Nitroglycerin*

11. Headache, dizziness and hypotension are side effects of Nitroglycerin?

a. True
b. False

12. King Airways require Base Station contact before insertion.

a. True
b. False

13. How much Aspirin is *given to the cardiac patient?*

a. 324mg (4 *tablets of81mg)*
b. 1296mg (4 *tablets of324mg*)
c. *500mg* (5 *tablets of81mg)*
d. l00mg (2 *tablets of81mg)*