

Interfacility/Interregional Ground Transport Policy

Purpose:

To provide consistent guidelines to Region 6 EMS agencies/providers and hospital personnel for interfacility/interregional transports.

Note:

This policy assumes that all EMS agencies/providers that provide interfacility/interregional transports have had System specific training for such transports.

Policy:

1. An attending physician, Emergency Department physician, or physician designee will authorize or request interfacility transports.
2. The transferring physician or physician designee will determine the appropriate receiving facility.
3. The transferring physician or physician designee will receive confirmation of acceptance of the patient from the receiving facility and the receiving physician or physician designee.
4. It is the responsibility of the transferring physician or physician designee to indicate what level of service and care is required for the transport based on the severity/complexity of the patient condition.
5. EMS agencies providing interfacility transports may only function to their level of licensure as defined by the National EMS Education Standards and Department regulations unless otherwise stated in this policy.
6. Any patient requiring care at a level higher than the highest level of prehospital care provider available must be transported with an RN or other appropriate professional personnel.
7. Prior to the transport, EMS providers must obtain written orders from the transferring physician or physician designee for all fluids and/or medications being transferred with the patient. EMS providers may only administer/monitor fluids and medications listed within this policy and the Region 6 protocols.

Online Medical Control:

Medical Control (MC) may be defined as either the EMS Medical Director, the transferring or receiving MD and as a last resort the ED physician of the transferring or receiving hospital.

In any situation that the EMS Provider needs to contact a physician for medical direction they will first attempt to contact the transferring MD or the receiving MD. If unable to reach either one, the EMS MD can be contacted. As a last resort, use on-line medical control at the sending or receiving facility. Any orders from on-line medical control will supersede written orders.

If the EMS Provider is unable to contact the receiving or sending facility, the EMS Provider will follow Region 6 EMS Protocols until contact can be established. In a situation when medical control is unreachable and intervention is necessary, the transport team will divert to the nearest appropriate medical facility.

Considerations for Transport:

Any Region 6 agency reserves the right to deny transport under the following conditions:

1. If providing the interfacility transport will impede the ability for the agency to provide 911 response within their response area due to staffing or equipment.
2. If it is deemed the patient is not stable enough for ground transport after consultation with the Medical Director or Medical Control.
3. If the safety of the patient and crew is at significant risk (i.e. weather, road conditions, violent patient, etc.).
4. Patients in active labor (when birth is imminent).
5. Active CPR in progress.

Requesting Additional Personnel:

When the EMS provider anticipates that they will require more assistance to appropriately care for the patient during transfer, they shall request the transferring physician/health care provider to provide appropriately trained hospital staff to accompany the patient and assist. The EMS provider must contact Medical Control for medical direction in all situations where they are not comfortable with the circumstances of the transfer. **The transfer will not occur unless the EMS provider and MC are confident the personnel and equipment are appropriate for transfer.**

Levels of EMS Interfacility Transports

Basic Life Support (BLS) interfacility transport

Minimum staffing: 2 EMT-Basic providers

Includes basic airway management, cardiopulmonary resuscitation including the use of AED's, basic shock management and control of bleeding, basic fracture management and medications within the Region 6 BLS protocols:

ASA
DuoNeb
Epinephrine 1:1000 (anaphylaxis only)
Naloxone
Nitroglycerin
Oral glucose

Basic providers may also transport patients with the following:

- Foley catheters
- Gastric devices (i.e., NG tubes, G tubes, ostomy equipment)
- Saline locks
- Wound drains
- Clamped Vascular devices (i.e., Central lines, Groshong catheters, PIC lines)*
***May not be accessed by Basic providers**

Intermediate Life Support (ILS) interfacility transport

Minimum staffing: 1 EMT-Intermediate and 1 EMT-Basic

Includes all BLS services, cardiac monitoring, IV cannulation/fluid therapy, advanced airway management and medications within the Region 6 ILS protocols:

Adenosine	Fentanyl
Amiodarone	Glucagon
Atropine	Hypertonic saline, 3.3%
Aspirin	Lidocaine
Dextrose, 50%	Methylprednisolone
Dextrose 10%	Midazolam
Diphenhydramine	Morphine
DuoNeb	Naloxone
Epinephrine 1:10,000	Nitroglycerin
Epinephrine 1:1000	Ondansetron
	Oral glucose

ILS providers may also transport patients with the following:

- CPAP / BiPAP
- IV infusion pumps

Advanced Life Support (ALS) interfacility transport

Minimum staffing: 1 EMT-Paramedic or Prehospital RN and 1 EMT-Basic

Includes all BLS and ILS services, cardiac monitoring (including cardiac pacing, manual defibrillation, and cardioversion) and administration/monitoring of medications within the Region 6 ALS protocols:

Adenosine	Glucagon
Amiodarone	Lidocaine
Atropine	Magnesium sulfate
Aspirin	Methylprednisolone
Dextrose 50%	Midazolam
Dextrose 10%	Morphine
Diltiazem	Naloxone
Diphenhydramine	Nitroglycerin
Dopamine	Ondansetron
DuoNeb	Oral glucose
Epinephrine 1:10,000	Sodium bicarbonate
Epinephrine 1:1000	Tranexamic acid
Fentanyl	

The following additional fluids and medications may also be transported by ALS providers:

All crystalloid and colloid solutions	Heparin drip
Acetadote	Hydralazine
Blood and Blood products (already initiated)	Hydrocortisone sodium succinate
IIb/IIIa glycoprotein inhibitors (Aggrastat,	Hydroxyzine
Reopro, Integrilin)	Isoproterenol
Antibiotics	Ketorolac
Atenolol	Labetalol (drip only)
Calcium chloride	Levophed
Calcium gluconate	Lorazepam
Cardene (drip only)	Mannitol
Dexamethasone sodium phosphate	Metoprolol (drip only)
Diazepam	Nifedipine (tabs)
Dobutamine	Nitroglycerine drip
Fentanyl drip	Oxytocin
Fosphenytoin	Octreotide

Phenobarbital (**drip only**)
Potassium (**no faster than 10 mEq/hr**)
Pralidoxime chloride
Propranolol (**drip only**)

Protonix
Racemic epinephrine
Sodium nitroprusside

ALS providers may also transport patients with the following:

- Pain medication pumps
- Femoral artery sheaths
- Chest tubes; with written physician orders. If mechanical suction, the amount of mechanical suction must be specified. Refer to "CHEST TUBE-POST INSERTION MANAGEMENT" in Care Guidelines.

One additional appropriately licensed healthcare provider in the patient compartment is required for the following:

All intubated patients (**Does not apply to stable ventilator dependent trach patients**)

****If not listed above or in the Region 6 protocols, a Registered Nurse is required to accompany the patient during transfer/transport.**

April 2005

Last Revised: November 2019