

Is your child up to date on immunizations?

Vaccines can prevent or help protect against deadly and difficult-to-treat diseases. And the shots don't end after infancy. Check below for the recommended ages of each immunization and determine if your child is current with his or her vaccines. Talk with your provider about an immunization schedule for your child.

Birth

- Recombivax HB (Hepatitis B)

2 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (Haemophilus influenzae type B)

4 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (Haemophilus influenzae type B)

6 months

- Pediarix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus/Hepatitis B)
- Prevnar 13 (Pneumococcal conjugate)
- RotaTeq (Rotavirus)
- Hiberix (Haemophilus influenzae type B)

12 months

- M-M-R II (Measles, Mumps, Rubella)
- Varivax (Varicella)
- Prevnar 13 (Pneumococcal conjugate)
- Havrix (Hepatitis A)

15 months

- Infanrix (Diphtheria, tetanus, and acellular pertussis [DTaP])
- Hiberix (Haemophilus influenzae type B)*

18 months

- Havrix (Hepatitis A)

4–6 years

- ProQuad (Measles, Mumps, Rubella/Varivax [Varicella]) or separate injections
- Kinrix (Diphtheria, tetanus, and acellular pertussis [DTaP]/Inactivated poliovirus)

11–12 years

- Boostrix (Tetanus, diphtheria, and acellular pertussis [Tdap])
- Menveo (Meningococcal Group A, C, Y)
- Gardasil 9 (Human papillomavirus) – 2 doses 6 months apart (over age 15 years requires 3 doses)

16 years

- Menveo (Meningococcal Group A, C, W, Y)
- Bexsero (Meningococcal Group B) – 2 doses**

Additional recommendations

- Influenza – Yearly for all children ages 6 months and up, if no contraindications
- COVID-19 – Children ages 5–17 may consider the two-dose Pfizer vaccine as well as a booster. (While the vaccine may not be available at the practice where your child is seen, the office can assist you in getting your child's vaccinations scheduled near you.)

PED-0126-OSF-04-23

NOTE: This schedule is for preventive vaccinations. For catch-up schedules or for patients with chronic conditions, see the CDC Immunization Schedule for vaccine recommendations.

*Can be administered as early as 12 months old

**For those ages 16-23 with no increased risk, discuss with provider.
Requires 2 doses, 1 month apart.



OSF[®]
HEALTHCARE